

B22

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 January 2003 (30.01.2003)


PCT

(10) International Publication Number
WO 03/008604 A2

- (51) International Patent Classification⁷: C12P 13/00 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (21) International Application Number: PCT/EP02/07352
- (22) International Filing Date: 3 July 2002 (03.07.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
101 35 051.1 18 July 2001 (18.07.2001) DE
60/306,867 23 July 2001 (23.07.2001) US
- (71) Applicant (*for all designated States except US*): DE-GUSSA AG [DE/DE]; Bennigsenplatz 1, 40474 Düsseldorf (DE).
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): HERMANN, Thomas [DE/DE]; Zirkonstrasse 8, 33739 Bielefeld (DE).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Declaration under Rule 4.17:
— of inventorship (Rule 4.17(iv)) for US only

[Continued on next page]

(54) Title: PROCESS FOR THE PREPARATION OF L-AMINO ACIDS USING STRAINS OF THE ENTEROBACTERIACEAE FAMILY WHICH CONTAIN AN ATTENUATED ACEB GENE



(57) Abstract: The invention relates to a process for the preparation of L-amino acids, in particular L-threonine, in which the following steps are carried out: a) fermentation of microorganisms of the Enterobacteriaceae family which produce the desired L-amino acid and in which the aceB gene, or the nucleotide sequence which codes for this, is attenuated, in particular eliminated, b) concentration of the L-amino acid in the medium or in the cells of the bacteria, and c) isolation of the L-amino acid.

WO 03/008604 A2



Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**Process for the Preparation of L-Amino Acids using
Strains of the Enterobacteriaceae Family which Contain
an Attenuated aceB Gene**

Field of the Invention

- 5 This invention relates to a process for the preparation of L-amino acids, in particular L-threonine, using strains of the Enterobacteriaceae family in which the aceB gene is attenuated.

Prior Art

- 10 L-Amino acids, in particular L-threonine, are used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and very particularly in animal nutrition.

- It is known to prepare L-amino acids by fermentation of strains of Enterobacteriaceae, in particular Escherichia coli (E. coli) and Serratia marcescens. Because of their great importance, work is constantly being undertaken to improve the preparation processes. Improvements to the process can relate to fermentation measures, such as e.g. stirring and supply of oxygen, or the composition of the nutrient media, such as e.g. the sugar concentration during the fermentation, or the working up to the product form, by e.g. ion exchange chromatography, or the intrinsic output properties of the microorganism itself.

- 25 Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to antimetabolites, such as e.g. the threonine analogue α -amino- β -hydroxyvaleric acid (AHV), or are auxotrophic for metabolites of regulatory importance and produce L-amino acid, such as e.g. L-threonine, are obtained in this manner.

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of strains of the Enterobacteriaceae family which produce L-amino acids, by amplifying individual amino acid biosynthesis genes and investigating the effect on the production.

Object of the Invention

The object of the invention is to provide new measures for improved fermentative preparation of L-amino acids, in particular L-threonine.

Summary of the Invention

The invention provides a process for the fermentative preparation of L-amino acids, in particular L-threonine, using microorganisms of the Enterobacteriaceae family which in particular already produce L-amino acids and in which the nucleotide sequence which codes for the aceB gene is attenuated.

Detailed Description of the Invention

Where L-amino acids or amino acids are mentioned in the following, this means one or more amino acids, including their salts, chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine. L-Threonine is particularly preferred.

The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by using a weak promoter or a gene or allele which codes for a corresponding enzyme with a low activity or inactivates the

corresponding enzyme (protein) or gene, and optionally combining these measures.

By attenuation measures, the activity or concentration of the corresponding protein is in general reduced to 0 to 75%, 0 to 50%, 0 to 25%, 0 to 10% or 0 to 5% of the activity or concentration of the wild-type protein or of the activity or concentration of the protein in the starting microorganism.

The process comprises carrying out the following steps:

- 10 a) fermentation of microorganisms of the Enterobacteriaceae family in which the aceB gene is attenuated,
 - b) concentration of the corresponding L-amino acid in the medium or in the cells of the microorganisms of the Enterobacteriaceae family, and
 - 15 c) isolation of the desired L-amino acid, constituents of the fermentation broth and/or the biomass in its entirety or portions (> 0 to 100 %) thereof optionally remaining in the product.
- 20 The microorganisms which the present invention provides can produce L-amino acids from glucose, sucrose, lactose, fructose, maltose, molasses, optionally starch, optionally cellulose or from glycerol and ethanol. They are representatives of the Enterobacteriaceae family chosen
- 25 from the genera *Escherichia*, *Erwinia*, *Providencia* and *Serratia*. The genera *Escherichia* and *Serratia* are preferred. Of the genus *Escherichia* the species *Escherichia coli* and of the genus *Serratia* the species *Serratia marcescens* are to be mentioned in particular.
- 30 Suitable strains, which produce L-threonine in particular, of the genus *Escherichia*, in particular of the species *Escherichia coli*, are, for example

- Escherichia coli TF427
Escherichia coli H4578
Escherichia coli KY10935
Escherichia coli VNIIGenetika MG442
5 Escherichia coli VNIIGenetika M1
Escherichia coli VNIIGenetika 472T23
Escherichia coli BKIIM B-3996
Escherichia coli kat 13
Escherichia coli KCCM-10132
- 10 Suitable L-threonine-producing strains of the genus
Serratia, in particular of the species Serratia marcescens,
are, for example
- Serratia marcescens HNr21
Serratia marcescens TLR156
15 Serratia marcescens T2000
- Strains from the Enterobacteriaceae family which produce L-
threonine preferably have, inter alia, one or more genetic
or phenotypic features chosen from the group consisting of:
resistance to α -amino- β -hydroxyvaleric acid, resistance to
20 thialysine, resistance to ethionine, resistance to α -
methylserine, resistance to diaminosuccinic acid,
resistance to α -aminobutyric acid, resistance to
borrelidin, resistance to rifampicin, resistance to valine
analogues, such as, for example, valine hydroxamate,
25 resistance to purine analogues, such as, for example, 6-
dimethylaminopurine, a need for L-methionine, optionally a
partial and compensable need for L-isoleucine, a need for
meso-diaminopimelic acid, auxotrophy in respect of
threonine-containing dipeptides, resistance to L-threonine,
30 resistance to L-homoserine, resistance to L-lysine,
resistance to L-methionine, resistance to L-glutamic acid,
resistance to L-aspartate, resistance to L-leucine,
resistance to L-phenylalanine, resistance to L-serine,
resistance to L-cysteine, resistance to L-valine,
35 sensitivity to fluoropyruvate, defective threonine

dehydrogenase, optionally an ability for sucrose utilization, enhancement of the threonine operon, enhancement of homoserine dehydrogenase I-aspartate kinase I, preferably of the feed back resistant form, enhancement
5 of homoserine kinase, enhancement of threonine synthase, enhancement of aspartate kinase, optionally of the feed back resistant form, enhancement of aspartate semialdehyde dehydrogenase, enhancement of phosphoenol pyruvate carboxylase, optionally of the feed back resistant form,
10 enhancement of phosphoenol pyruvate synthase, enhancement of transhydrogenase, enhancement of the RhtB gene product, enhancement of the RhtC gene product, enhancement of the YfiK gene product, enhancement of a pyruvate carboxylase, and attenuation of acetic acid formation.

15 It has been found that microorganisms of the Enterobacteriaceae family produce L-amino acids, in particular L-threonine, in an improved manner after attenuation, in particular elimination, of the aceB gene

The nucleotide sequences of the genes of Escherichia coli
20 belong to the prior art and can also be found in the genome sequence of Escherichia coli published by Blattner et al. (Science 277: 1453 - 1462 (1997)).

The aceB gene is described, inter alia, by the following data:

25 Description: Malate synthase A
EC No.: 4.1.3.2
Reference: Byrne et al.; Nucleic Acids Research
16(19), 9342 (1988); Byrne et al.; Nucleic
Acids Research 16(22), 10924 (1988); Cortay
30 et al.; Biochimie 71(9-10): 1043-9 (1989)
Accession No.: AE000474
Alternative gene name: mas

The nucleic acid sequences can be found in the databanks of the National Center for Biotechnology Information (NCBI) of the National Library of Medicine (Bethesda, MD, USA), the nucleotide sequence databank of the European Molecular
5 Biologies Laboratories (EMBL, Heidelberg, Germany or Cambridge, UK) or the DNA databank of Japan (DDBJ, Mishima, Japan).

The genes described in the text references mentioned can be used according to the invention. Alleles of the genes which
10 result from the degeneracy of the genetic code or due to "sense mutations" of neutral function can furthermore be used.

To achieve an attenuation, for example, expression of the gene or the catalytic properties of the enzyme proteins can
15 be reduced or eliminated. The two measures can optionally be combined.

The reduction in gene expression can take place by suitable culturing, by genetic modification (mutation) of the signal structures of gene expression or also by the antisense-RNA
20 technique. Signal structures of gene expression are, for example, repressor genes, activator genes, operators, promoters, attenuators, ribosome binding sites, the start codon and terminators. The expert can find information in this respect, inter alia, for example, in Jensen and Hammer
25 (Biotechnology and Bioengineering 58: 191-195 (1998)), in Carrier and Keasling (Biotechnology Progress 15: 58-64 (1999), Franch and Gerdes (Current Opinion in Microbiology 3: 159-164 (2000)) and in known textbooks of genetics and molecular biology, such as, for example, the textbook of
30 Knippers ("Molekulare Genetik [Molecular Genetics]", 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995) or that of Winnacker ("Gene und Klone [Genes and Clones]", VCH Verlagsgesellschaft, Weinheim, Germany, 1990).

Mutations which lead to a change or reduction in the catalytic properties of enzyme proteins are known from the prior art. Examples which may be mentioned are the works of Qiu and Goodman (Journal of Biological Chemistry 272: 8611-8617 (1997)), Yano et al. (Proceedings of the National Academy of Sciences of the United States of America 95: 5511-5515 (1998)), Wentz and Schachmann (Journal of Biological Chemistry 266: 20833-20839 (1991)). Summarizing descriptions can be found in known textbooks of genetics and molecular biology, such as e.g. that by Hagemann ("Allgemeine Genetik [General Genetics]", Gustav Fischer Verlag, Stuttgart, 1986).

Possible mutations are transitions, transversions, insertions and deletions. Depending on the effect of the amino acid exchange on the enzyme activity, "missense mutations" or "nonsense mutations" are referred to. Insertions or deletions of at least one base pair in a gene lead to "frame shift mutations", which lead to incorrect amino acids being incorporated or translation being interrupted prematurely. If a stop codon is formed in the coding region as a consequence of the mutation, this also leads to a premature termination of the translation. Deletions of several codons typically lead to a complete loss of the enzyme activity. Instructions on generation of such mutations are prior art and can be found in known textbooks of genetics and molecular biology, such as e.g. the textbook by Knippers ("Molekulare Genetik [Molecular Genetics]", 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995), that by Winnacker ("Gene und Klone [Genes and Clones]", VCH Verlagsgesellschaft, Weinheim, Germany, 1990) or that by Hagemann ("Allgemeine Genetik [General Genetics]", Gustav Fischer Verlag, Stuttgart, 1986).

Suitable mutations in the genes, such as, for example, deletion mutations, can be incorporated into suitable strains by gene or allele replacement.

A conventional method is the method, described by Hamilton et al. (Journal of Bacteriology 171: 4617-4622 (1989)), of gene replacement with the aid of a conditionally replicating pSC101 derivative pMAK705. Other methods
5 described in the prior art, such as, for example, those of Martinez-Morales et al. (Journal of Bacteriology 181: 7143-7148 (1999)) or those of Boyd et al. (Journal of Bacteriology 182: 842-847 (2000)), can likewise be used.

It is also possible to transfer mutations in the particular
10 genes or mutations which affect expression of the particular genes into various strains by conjugation or transduction.

It may furthermore be advantageous for the production of L-amino acids, in particular L-threonine, with strains of the
15 Enterobacteriaceae family, in addition to attenuation of the aceB gene, for one or more enzymes of the known threonine biosynthesis pathway or enzymes of anaplerotic metabolism or enzymes for the production of reduced nicotinamide adenine dinucleotide phosphate or enzymes of
20 glycolysis or PTS enzymes or enzymes of sulfur metabolism to be enhanced.

The term "enhancement" in this connection describes the increase in the intracellular activity of one or more enzymes or proteins in a microorganism which are coded by
25 the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or a gene which codes for a corresponding enzyme or protein with a high activity, and optionally combining these measures.

30 By enhancement measures, in particular over-expression, the activity or concentration of the corresponding protein is in general increased by at least 10%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400% or 500%, up to a maximum of 1000% or 2000%, based on that of the wild-type protein or the

activity or concentration of the protein in the starting microorganism.

Thus, for example, at the same time one or more of the genes chosen from the group consisting of

- 5 • the thrABC operon which codes for aspartate kinase, homoserine dehydrogenase, homoserine kinase and threonine synthase (US-A-4,278,765),
- the pyc gene of Corynebacterium glutamicum which codes for pyruvate carboxylase (WO 99/18228),
- 10 • the pps gene which codes for phosphoenol pyruvate synthase (Molecular and General Genetics 231(2): 332-336 (1992)),
- the ppc gene which codes for phosphoenol pyruvate carboxylase (Gene 31: 279-283 (1984)),
- 15 • the pntA and pntB genes which code for transhydrogenase (European Journal of Biochemistry 158: 647-653 (1986)),
- the rhtB gene which imparts homoserine resistance (EP-A-0 994 190),
- the mqo gene which codes for malate:quinone
20 oxidoreductase (WO 02/06459),
- the rhtC gene which imparts threonine resistance (EP-A-1 013 765),
- the thrE gene of Corynebacterium glutamicum which codes for the threonine export protein (WO 01/92545),
- 25 • the gdhA gene which codes for glutamate dehydrogenase (Nucleic Acids Research 11: 5257-5266 (1983); Gene 23: 199-209 (1983)),

- the hns gene which codes for the DNA-binding protein HLP-II (Molecular and General Genetics 212: 199-202 (1988)),
- 5 • the pgm gene which codes for phosphoglucomutase (Journal of Bacteriology 176: 5847-5851 (1994)),
- the fba gene which codes for fructose biphosphate aldolase (Biochemical Journal 257: 529-534 (1989)),
- 10 • the ptsH gene of the ptsHIcrr operon which codes for the phosphohistidine protein hexose phosphotransferase of the phosphotransferase system PTS (Journal of Biological Chemistry 262: 16241-16253 (1987)),
- the ptsI gene of the ptsHIcrr operon which codes for enzyme I of the phosphotransferase system PTS (Journal of Biological Chemistry 262: 16241-16253 (1987)),
- 15 • the crr gene of the ptsHIcrr operon which codes for the glucose-specific IIA component of the phosphotransferase system PTS (Journal of Biological Chemistry 262: 16241-16253 (1987)),
- the ptsG gene which codes for the glucose-specific IIBC component (Journal of Biological Chemistry 261: 16398-16403 (1986)),
- 20 • the lrp gene which codes for the regulator of the leucine regulon (Journal of Biological Chemistry 266: 10768-10774 (1991)),
- 25 • the mopB gene which codes for 10 Kd chaperone (Journal of Biological Chemistry 261: 12414-12419 (1986)) and is also known by the name groES,
- the ahpC gene of the ahpCF operon which codes for the small sub-unit of alkyl hydroperoxide reductase
- 30 (Proceedings of the National Academy of Sciences of the United States of America 92: 7617-7621 (1995)),

- the *ahpF* gene of the *ahpCF* operon which codes for the large sub-unit of alkyl hydroperoxide reductase (Proceedings of the National Academy of Sciences of the United States of America 92: 7617-7621 (1995)),
- 5 • the *cysK* gene which codes for cysteine synthase A (Journal of Bacteriology 170: 3150-3157 (1988)),
- the *cysB* gene which codes for the regulator of the *cys* regulon (Journal of Biological Chemistry 262: 5999-6005 (1987)),
- 10 • the *cysJ* gene of the *cysJIH* operon which codes for the flavoprotein of NADPH sulfite reductase (Journal of Biological Chemistry 264: 15796-15808 (1989), Journal of Biological Chemistry 264: 15726-15737 (1989)),
- the *cysI* gene of the *cysJIH* operon which codes for the
15 haemoprotein of NADPH sulfite reductase (Journal of Biological Chemistry 264: 15796-15808 (1989), Journal of Biological Chemistry 264: 15726-15737 (1989)) and
- the *cysH* gene of the *cysJIH* operon which codes for
20 adenylyl sulfate reductase (Journal of Biological Chemistry 264: 15796-15808 (1989), Journal of Biological Chemistry 264: 15726-15737 (1989))

can be enhanced, in particular over-expressed.

The use of endogenous genes is in general preferred.

- "Endogenous genes" or "endogenous nucleotide sequences" are
25 understood as meaning the genes or nucleotide sequences present in the population of a species.

It may furthermore be advantageous for the production of L-amino acids, in particular L-threonine, in addition to attenuation of the *aceB* gene, for one or more of the genes
30 chosen from the group consisting of

- the tdh gene which codes for threonine dehydrogenase (Journal of Bacteriology 169: 4716-4721 (1987)),
- the mdh gene which codes for malate dehydrogenase (E.C. 1.1.1.37) (Archives in Microbiology 149: 36-42 (1987)),
- 5 • the gene product of the open reading frame (orf) yjfa (Accession Number AAC77180 of the National Center for Biotechnology Information (NCBI, Bethesda, MD, USA)),
- the gene product of the open reading frame (orf) ytfP (Accession Number AAC77179 of the National Center for
10 Biotechnology Information (NCBI, Bethesda, MD, USA)),
- the pckA gene which codes for the enzyme phosphoenol pyruvate carboxykinase (Journal of Bacteriology 172: 7151-7156 (1990)),
- the poxB gene which codes for pyruvate oxidase (Nucleic
15 Acids Research 14(13): 5449-5460 (1986)),
- the aceA gene which codes for the enzyme isocitrate lyase (Journal of Bacteriology 170: 4528-4536 (1988)),
- the dgsA gene which codes for the DgsA regulator of the phosphotransferase system (Bioscience, Biotechnology and
20 Biochemistry 59: 256-261 (1995)) and is also known under the name of the mlc gene,
- the fruR gene which codes for the fructose repressor (Molecular and General Genetics 226: 332-336 (1991)) and is also known under the name of the cra gene,
- 25 • the rpoS gene which codes for the sigma³⁸ factor (WO 01/05939) and is also known under the name of the katF gene,
- the aspA gene which codes for aspartate ammonium lyase (Nucleic Acids Research 13(6): 2063-2074 (1985)),

- the aceK gene which codes for isocitrate dehydrogenase kinase/phosphatase (Journal of Bacteriology 170(1): 89-97 (1988)) and
- the ugpB gene which codes for the periplasmic binding protein of the sn-glycerol 3-phosphate transport system (Molecular Microbiology 2(6): 767-775 (1988))

to be attenuated, in particular eliminated or for the expression thereof to be reduced.

It may furthermore be advantageous for the production of L-amino acids, in particular L-threonine, in addition to attenuation of the aceB gene, to eliminate undesirable side reactions (Nakayama: "Breeding of Amino Acid Producing Microorganisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

The microorganisms produced according to the invention can be cultured in the batch process (batch culture), the fed batch process (feed process) or the repeated fed batch process (repetitive feed process). A summary of known culture methods is described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik [Bioprocess Technology 1. Introduction to Bioprocess Technology (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen [Bioreactors and Peripheral Equipment] (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981).

Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and optionally cellulose, oils and fats, such as e.g. soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as
5 e.g. palmitic acid, stearic acid and linoleic acid, alcohols, such as e.g. glycerol and ethanol, and organic acids, such as e.g. acetic acid, can be used as the source of carbon. These substances can be used individually or as a mixture.

- 10 Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be
15 used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus.

- 20 The culture medium must furthermore comprise salts of metals, such as e.g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the abovementioned substances.
25 Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

- Basic compounds, such as sodium hydroxide, potassium
30 hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture. Antifoams, such as e.g. fatty acid polyglycol esters, can be employed to control the development of foam. Suitable
35 substances having a selective action, e.g. antibiotics, can

be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as e.g. air, are introduced into the culture. The temperature of the
5 culture is usually 25°C to 45°C, and preferably 30°C to 40°C. Culturing is continued until a maximum of L-amino acids or L-threonine has formed. This target is usually reached within 10 hours to 160 hours.

The analysis of L-amino acids can be carried out by anion
10 exchange chromatography with subsequent ninhydrin derivation, as described by Spackman et al. (Analytical Chemistry 30: 1190-1206 (1958)), or it can take place by reversed phase HPLC as described by Lindroth et al. (Analytical Chemistry 51: 1167-1174 (1979)).

15 The process according to the invention is used for the fermentative preparation of L-amino acids, such as; for example, L-threonine, L-isoleucine, L-valine, L-methionine, L-homoserine and L-lysine, in particular L-threonine.

A pure culture of the Escherichia coli K-12 strain
20 DH5 α /pMAK705 was deposited as DSM 13720 on 8th September 2000 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty.

25 The present invention is explained in more detail in the following with the aid of embodiment examples.

The isolation of plasmid DNA from Escherichia coli and all techniques of restriction, ligation, Klenow and alkaline phosphatase treatment are carried out by the method of
30 Sambrook et al. (Molecular Cloning - A Laboratory Manual (1989) Cold Spring Harbor Laboratory Press). Unless described otherwise, the transformation of Escherichia coli is carried out by the method of Chung et al. (Proceedings

of the National Academy of Sciences of the United States of America 86: 2172-2175 (1989)).

The incubation temperature for the preparation of strains and transformants is 37°C. Temperatures of 30°C and 44°C
5 are used in the gene replacement method of Hamilton et al.

Example 1

Construction of the deletion mutation of the aceB gene

Parts of the gene regions lying upstream and downstream of the aceB gene and parts of the 5' and 3' region of the aceB
10 gene are amplified from Escherichia coli K12 using the polymerase chain reaction (PCR) and synthetic oligonucleotides. Starting from the nucleotide sequence of the aceB gene and sequences lying upstream and downstream in E. coli K12 MG1655 (SEQ ID No. 1, Accession Number
15 AE000474), the following PCR primers are synthesized (MWG Biotech, Ebersberg, Germany):

aceB5'-1: 5' - TTCGGATCCATGACGAGGAG - 3' (SEQ ID No. 3)

aceB5'-2: 5' - TTGCCAACAGTGCCTGATAG - 3' (SEQ ID No. 4)

aceB3'-1: 5' - ATGCTTACTCACGCCTGTTG - 3' (SEQ ID No. 5)

20 aceB3'-2: 5' - CATGTGCAGATGCTCCATAG - 3' (SEQ ID No. 6)

The chromosomal E. coli K12 MG1655 DNA employed for the PCR is isolated according to the manufacturer's instructions with "Qiagen Genomic-tips 100/G" (QIAGEN, Hilden, Germany). A DNA fragment approx. 650 bp in size from the 5' region of
25 the aceB gene region (called aceB5') and a DNA fragment approx. 700 bp in size from the 3' region of the aceB gene region (called aceB3') can be amplified with the specific primers under standard PCR conditions (Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications,
30 Academic Press) with Taq-DNA polymerase (Gibco-BRL, Eggenstein, Germany). The PCR products are each ligated

with the vector pCRII-TOPO (TOPO TA Cloning Kit, Invitrogen, Groningen, The Netherlands) in accordance with the manufacturer's instructions and transformed into the E. coli strain TOP10F'. Selection of plasmid-carrying cells
5 takes place on LB agar, to which 50 µg/ml ampicillin are added. After isolation of the plasmid DNA, the vector pCRII-TOPOaceB3' is cleaved with the restriction enzymes XbaI and Ecl136II. The aceB5' fragment is isolated after separation in 0.8% agarose gel with the aid of the QIAquick
10 Gel Extraction Kit (QIAGEN, Hilden, Germany). After isolation of the plasmid DNA the vector pCRII-TOPOaceB5' is cleaved with the enzymes EcoRV and XbaI and ligated with the aceB5' fragment isolated. The E. coli strain DH5α is transformed with the ligation batch and plasmid-carrying
15 cells are selected on LB agar, to which 50 µg/ml ampicillin are added. After isolation of the plasmid DNA those plasmids in which the mutagenic DNA sequence shown in SEQ ID No. 7 is cloned are detected by control cleavage with the enzymes BclI, HincII, SpeI and SphI. One of the
20 plasmids is called pCRII-TOPOΔaceB (=pCRII-TOPOdeltaaceB).

Example 2

Construction of the replacement vector pMAK705ΔaceB

The ΔaceB allele described in example 1 is isolated from the vector pCRII-TOPOΔaceB after restriction with the
25 enzymes KpnI and XbaI and separation in 0.8% agarose gel, and ligated with the plasmid pMAK705 (Hamilton et al., Journal of Bacteriology 171: 4617-4622 (1989)), which has been digested with the enzymes KpnI and XbaI. The ligation batch is transformed in DH5α and plasmid-carrying cells are
30 selected on LB agar, to which 20 µg/ml chloramphenicol are added. Successful cloning is demonstrated after isolation of the plasmid DNA and cleavage with the enzymes BamHI, ClaI, PvuI and XbaI. The replacement vector formed, pMAK705ΔaceB (= pMAK705deltaaceB), is shown in Figure 1.

Example 3

Position-specific mutagenesis of the aceB gene in the E. coli strain MG442

The L-threonine-producing E. coli strain MG442 is described
5 in the patent specification US-A- 4,278,765 and deposited
as CMIM B-1628 at the Russian National Collection for
Industrial Microorganisms (VKPM, Moscow, Russia).

For replacement of the chromosomal aceB gene with the
plasmid-coded deletion construct, MG442 is transformed with
10 the plasmid pMAK705 Δ aceB. The gene replacement is carried
out using the selection method described by Hamilton et al.
(Journal of Bacteriology 171: 4617-4622 (1989)) and is
verified by standard PCR methods (Innis et al. (1990) PCR
Protocols. A Guide to Methods and Applications, Academic
15 Press) with the following oligonucleotide primers:

aceB5'-1: 5' - TTCGGATCCATGACGAGGAG - 3' (SEQ ID No. 3)

aceB3'-2: 5' - CATGTGCAGATGCTCCATAG - 3' (SEQ ID No. 6)

After replacement has taken place, MG442 contains the form
of the Δ aceB allele shown in SEQ ID No. 8. The strain
20 obtained is called MG442 Δ aceB.

Example 4

Preparation of L-threonine with the strain MG442 Δ aceB

MG442 Δ aceB is multiplied on minimal medium with the
following composition: 3.5 g/l Na₂HPO₄*2H₂O, 1.5 g/l KH₂PO₄,
25 1 g/l NH₄Cl, 0.1 g/l MgSO₄*7H₂O, 2 g/l glucose, 20 g/l agar.
The formation of L-threonine is checked in batch cultures
of 10 ml contained in 100 ml conical flasks. For this,
10 ml of preculture medium of the following composition:
2 g/l yeast extract, 10 g/l (NH₄)₂SO₄, 1 g/l KH₂PO₄, 0.5 g/l
30 MgSO₄*7H₂O, 15 g/l CaCO₃, 20 g/l glucose are inoculated and
the batch is incubated for 16 hours at 37°C and 180 rpm on

- an ESR incubator from Kühner AG (Birsfelden, Switzerland). 250 µl of this preculture are transinoculated into 10 ml of production medium (25 g/l $(\text{NH}_4)_2\text{SO}_4$, 2 g/l KH_2PO_4 , 1 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.03 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.018 g/l $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 30 g/l CaCO_3 , 20 g/l glucose) and the batch is incubated for 48 hours at 37°C. After the incubation the optical density (OD) of the culture suspension is determined with an LP2W photometer from Dr. Lange (Düsseldorf, Germany) at a measurement wavelength of 660 nm.
- 10 The concentration of L-threonine formed is then determined in the sterile-filtered culture supernatant with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column reaction with ninhydrin detection.
- 15 The result of the experiment is shown in Table 1.

Table 1

Strain	OD (660 nm)	L-Threonine g/l
MG442	6.0	1.5
MG442 Δ aceB	4.8	2.1

Brief Description of the Figure:

- Figure 1: pMAK705 Δ aceB (= pMAK705deltaaceB)
- 20 The length data are to be understood as approx. data. The abbreviations and designations used have the following meaning:
- cat: Chloramphenicol resistance gene
 - rep-ts: Temperature-sensitive replication region of the plasmid pSC101
- 25

- aceB5': Part of the 5' region of the aceB gene and the region lying upstream
- aceB3': Part of the 3' region of the aceB gene and the region lying downstream

5 The abbreviations for the restriction enzymes have the following meaning

- BamHI: Restriction endonuclease from *Bacillus amyloliquefaciens* H
- ClaI: Restriction endonuclease from *Caryophanon latum*
- 10 • KpnI: Restriction endonuclease from *Klebsiella pneumoniae*
- PstI: Restriction endonuclease from *Paracoccus alcaliphilus*
- XbaI: Restriction endonuclease from *Xanthomonas badrii*

What is claimed is:

1. A process for the preparation of L-amino acids, in particular L-threonine, which comprises carrying out the following steps:
 - 5 a) fermentation of microorganisms of the Enterobacteriaceae family which produce the desired L-amino acid and in which the aceB gene, or the nucleotide sequence which codes for this, is attenuated, in particular eliminated,
 - 10 b) concentration of the desired L-amino acid in the medium or in the cells of the microorganisms, and
 - c) isolation of the desired L-amino acid, constituents of the fermentation broth and/or the biomass in its entirety or portions (> 0 to 100 %) thereof
 - 15 optionally remaining in the product.
2. A process as claimed in claim 1, wherein microorganisms in which further genes of the biosynthesis pathway of the desired L-amino acid are additionally enhanced are employed.
- 20 3. A process as claimed in claim 1, wherein microorganisms in which the metabolic pathways which reduce the formation of the desired L-amino acid are at least partly eliminated are employed.
4. A process as claimed in claim 1, wherein the expression
25 of the polynucleotide which codes for the aceB gene is attenuated, in particular eliminated.
5. A process as claimed in claim 1, wherein the regulatory and/or catalytic properties of the polypeptide (enzyme protein) for which the polynucleotide aceB codes are
30 reduced.

6. A process as claimed in claim 1, wherein, for the preparation of L-amino acids, microorganisms of the Enterobacteriaceae family in which in addition at the same time one or more of the genes chosen from the group consisting of:
- 5
- 6.1 the thrABC operon which codes for aspartate kinase, homoserine dehydrogenase, homoserine kinase and threonine synthase,
 - 6.2 the pyc gene which codes for pyruvate carboxylase,
 - 10 6.3 the pps gene which codes for phosphoenol pyruvate synthase,
 - 6.4 the ppc gene which codes for phosphoenol pyruvate carboxylase,
 - 15 6.5 the pntA and pntB genes which code for transhydrogenase,
 - 6.6 the rhtB gene which imparts homoserine resistance,
 - 6.7 the mgo gene which codes for malate:quinone oxidoreductase,
 - 20 6.8 the rhtC gene which imparts threonine resistance,
 - 6.9 the thrE gene which codes for the threonine export protein,
 - 25 6.10 the gdhA gene which codes for glutamate dehydrogenase,
 - 6.11 the hns gene which codes for the DNA-binding protein HLP-II,

- 6.12 the pgm gene which codes for phosphoglucomutase,
- 6.13 the fba gene which codes for fructose biphosphate aldolase,
- 5 6.14 the ptsH gene which codes for the phosphohistidine protein hexose phosphotransferase,
- 6.15 the ptsI gene which codes for enzyme I of the phosphotransferase system,
- 10 6.16 the crr gene which codes for the glucose-specific IIA component,
- 6.17 the ptsG gene which codes for the glucose-specific IIBC component,
- 15 6.18 the lrp gene which codes for the regulator of the leucine regulon,
- 6.19 the mopB gene which codes for 10 Kd chaperone,
- 6.20 the ahpC gene which codes for the small sub-unit of alkyl hydroperoxide reductase,
- 20 6.21 the ahpF gene which codes for the large sub-unit of alkyl hydroperoxide reductase,
- 6.22 the cysK gene which codes for cysteine synthase A,
- 6.23 the cysB gene which codes for the regulator of the cys regulon,
- 25 6.24 the cysJ gene which codes for the flavoprotein of NADPH sulfite reductase,
- 6.25 the cysI gene which codes for the haemoprotein of NADPH sulfite reductase and

6.26 the cysH gene which codes for adenylyl sulfate reductase,


is or are enhanced, in particular over-expressed, are fermented.

- 5 7. A process as claimed in claim 1, wherein, for the preparation of L-amino acids, microorganisms of the Enterobacteriaceae family in which in addition at the same time one or more of the genes chosen from the group consisting of:
- 10 7.1 the tdh gene which codes for threonine dehydrogenase,
- 7.2 the mdh gene which codes for malate dehydrogenase,
- 15 7.3 the gene product of the open reading frame (orf) yjfA,
- 7.4 the gene product of the open reading frame (orf) ytfP,
- 7.5 the pckA gene which codes for phosphoenolpyruvate carboxykinase
- 20 7.6 the poxB gene which codes for pyruvate oxidase
- 7.7 the aceA gene which codes for isocitrate lyase,
- 7.8 the dgsA gene which codes for the DgsA regulator of the phosphotransferase system,
- 25 7.9 the fruR gene which codes for the fructose repressor,
- 7.10 the rpoS gene which codes for the sigma³⁸ factor,

- 7.11 the aspA gene which codes for aspartate ammonium lyase,
- 7.12 the aceK gene which codes for isocitrate dehydrogenase kinase/phosphatase and
- 5 7.13 the ugpB gene which codes for the periplasmic binding protein of the sn-glycerol 3-phosphate transport system

is or are attenuated, in particular eliminated or reduced in expression, are fermented.

Figure 1:



SEQUENCE PROTOCOL

5 <110> Degussa AG
 <120> Process for the preparation of L-amino acids using strains of the Enterobacteriaceae family which contain an attenuated aceB gene
 10 <130> 020271 BT
 <160> 8
 15 <170> PatentIn version 3.1
 <210> 1
 <211> 1661
 <212> DNA
 <213> Escherichia coli
 20 <220>
 <221> CDS
 <222> (28)..(1629)
 <223> aceB gene.
 25 <400> 1
 ttccggtacca tgacgaggag ctgcacg atg act gaa cag gca aca aca acc gat 54
 Met Thr Glu Gln Ala Thr Thr Thr Asp
 1 5
 30 gaa ctg gct ttc aca agg ccg tat ggc gag cag gag aag caa att ctt 102
 Glu Leu Ala Phe Thr Arg Pro Tyr Gly Glu Gln Glu Lys Gln Ile Leu
 10 15 20 25
 35 act gcc gaa gcg gta gaa ttt ctg act gag ctg gtg acg cat ttt acg 150
 Thr Ala Glu Ala Val Glu Phe Leu Thr Glu Leu Val Thr His Phe Thr
 30 35 40
 40 cca caa cgc aat aaa ctt ctg gca gcg cgc att cag cag cag caa gat 198
 Pro Gln Arg Asn Lys Leu Leu Ala Ala Arg Ile Gln Gln Gln Gln Asp
 45 50 55
 att gat aac gga acg ttg cct gat ttt att tcg gaa aca gct tcc att 246
 Ile Asp Asn Gly Thr Leu Pro Asp Phe Ile Ser Glu Thr Ala Ser Ile
 45 60 65 70
 cgc gat gct gat tgg aaa att cgc ggg att cct gcg gac tta gaa gac 294
 Arg Asp Ala Asp Trp Lys Ile Arg Gly Ile Pro Ala Asp Leu Glu Asp
 75 80 85
 50 cgc cgc gta gag ata act ggc ccg gta gag cgc aag atg gtg atc aac 342
 Arg Arg Val Glu Ile Thr Gly Pro Val Glu Arg Lys Met Val Ile Asn
 90 95 100 105
 55 gcg ctc aac gcc aat gtg aaa gtc ttt atg gcc gat ttc gaa gat tca 390
 Ala Leu Asn Ala Asn Val Lys Val Phe Met Ala Asp Phe Glu Asp Ser
 110 115 120
 60 ctg gca cca gac tgg aac aaa gtg atc gac ggg caa att aac ctg cgt 438
 Leu Ala Pro Asp Trp Asn Lys Val Ile Asp Gly Gln Ile Asn Leu Arg
 125 130 135
 65 gaa tgg gtc aac ggc aac aac agc aac aac aac gaa gaa ggc aac att 436
 Asp Ala Val Asn Gly Thr Ile Ser Tyr Thr Asn Glu Ala Gly Lys Ile
 140 145 150

5	tac cag ctc aag ccc aat cca gcg gtt ttg att tgt cgg gta cgc ggt Tyr Gln Leu Lys Pro Asn Pro Ala Val Leu Ile Cys Arg Val Arg Gly 155 160 165	534
10	ctg cac ttg ccg gaa aaa cat gtc acc tgg cgt ggt gag gca atc ccc Leu His Leu Pro Glu Lys His Val Thr Trp Arg Gly Glu Ala Ile Pro 170 175 180 185	582
15	ggc agc ctg ttt gat ttt gcg ctc tat ttc ttc cac aac tat cag gca Gly Ser Leu Phe Asp Phe Ala Leu Tyr Phe Phe His Asn Tyr Gln Ala 190 195 200	630
20	ctg ttg gca aag ggc agt ggt ccc tat ttc tat ctg ccg aaa acc cag Leu Leu Ala Lys Gly Ser Gly Pro Tyr Phe Tyr Leu Pro Lys Thr Gln 205 210 215	678
25	tcc tgg cag gaa gcg gcc tgg tgg agc gaa gtc ttc agc tat gca gaa Ser Trp Gln Glu Ala Ala Trp Trp Ser Glu Val Phe Ser Tyr Ala Glu 220 225 230	726
30	gat cgc ttt aat ctg ccg gcg ggc acc atc aag gcg acg ttg ctg att Asp Arg Phe Asn Leu Pro Arg Gly Thr Ile Lys Ala Thr Leu Leu Ile 235 240 245	774
35	gaa acg ctg ccc gcc gtg ttc cag atg gat gaa atc ctt cac gcg ctg Glu Thr Leu Pro Ala Val Phe Gln Met Asp Glu Ile Leu His Ala Leu 250 255 260 265	822
40	cgt gac cat att gtt ggt ctg aac tgc ggt cgt tgg gat tac atc ttc Arg Asp His Ile Val Gly Leu Asn Cys Gly Arg Trp Asp Tyr Ile Phe 270 275 280	870
45	agc tat atc aaa acg ttg aaa aac tat ccc gat cgc gtc ctg cca gac Ser Tyr Ile Lys Thr Leu Lys Asn Tyr Pro Asp Arg Val Leu Pro Asp 285 290 295	918
50	aga cag gca gtg acg atg gat aaa cca ttc ctg aat gct tac tca cgc Arg Gln Ala Val Thr Met Asp Lys Pro Phe Leu Asn Ala Tyr Ser Arg 300 305 310	966
55	ctg ttg att aaa acc tgc cat aaa cgc ggt gct ttt gcg atg ggc ggc Leu Leu Ile Lys Thr Cys His Lys Arg Gly Ala Phe Ala Met Gly Gly 315 320 325	1014
60	atg gcg gcg ttt att ccg agc aaa gat gaa gag cac aat aac cag gtg Met Ala Ala Phe Ile Pro Ser Lys Asp Glu Glu His Asn Asn Gln Val 330 335 340 345	1062
65	ctc aac aaa gta aaa gcg gat aaa tcg ctg gaa gcc aat aac ggt cac Leu Asn Lys Val Lys Ala Asp Lys Ser Leu Glu Ala Asn Asn Gly His 350 355 360	1110
70	gat ggc aca tgg atc gct cac cca ggc ctt gcg gac acg gca atg gcg Asp Gly Thr Trp Ile Ala His Pro Gly Leu Ala Asp Thr Ala Met Ala 365 370 375	1158
75	gta ttc aac gac att ctc ggc tcc cgt aaa aat cag ctt gaa gtg atg Val Phe Asn Asp Ile Leu Gly Ser Arg Lys Asn Gln Leu Glu Val Met 380 385 390	1206
80	cgc gaa caa gac gcg ccg att act gcc gat cag ctg ctg gca cct tgt Arg Glu Gln Asp Ala Pro Ile Thr Ala Asp Gln Leu Leu Ala Pro Cys 395 400 405	1254

	gat ggt gaa cgc acc gaa gaa ggt atg cgc gcc aac att cgc gtg gct Asp Gly Glu Arg Thr Glu Glu Gly Met Arg Ala Asn Ile Arg Val Ala 410 415 420 425	1302
5	gtg cag tac atc gaa gcg tgg atc tct ggc aac ggc tgt gtg ccg att Val Gln Tyr Ile Glu Ala Trp Ile Ser Gly Asn Gly Cys Val Pro Ile 430 435 440	1350
10	tat ggc ctg atg gaa gat gcg gcg acg gct gaa att tcc cgt acc tcg Tyr Gly Leu Met Glu Asp Ala Ala Thr Ala Glu Ile Ser Arg Thr Ser 445 450 455	1398
15	atc tgg cag tgg atc cat cat caa aaa acg ttg agc aat ggc aaa ccg Ile Trp Gln Trp Ile His His Gln Lys Thr Leu Ser Asn Gly Lys Pro 460 465 470	1446
20	gtg acc aaa gcc ttg ttc cgc cag atg ctg ggc gaa gag atg aaa gtc Val Thr Lys Ala Leu Phe Arg Gln Met Leu Gly Glu Glu Met Lys Val 475 480 485	1494
25	att gcc agc gaa ctg ggc gaa gaa cgt ttc tcc cag ggg cgt ttt gac Ile Ala Ser Glu Leu Gly Glu Glu Arg Phe Ser Gln Gly Arg Phe Asp 490 495 500 505	1542
30	gat gcc gca cgc ttg atg gaa cag atc acc act tcc gat gag tta att Asp Ala Ala Arg Leu Met Glu Gln Ile Thr Thr Ser Asp Glu Leu Ile 510 515 520	1590
35	gat ttc ctg acc ctg cca ggc tac cgc ctg tta gcg taa accaccacat Asp Phe Leu Thr Leu Pro Gly Tyr Arg Leu Leu Ala 525 530	1639
40	aactatggag catctgcaca tg	1661
45	<210> 2 <211> 533 <212> PRT <213> Escherichia coli	
50	<400> 2 Met Thr Glu Gln Ala Thr Thr Thr Asp Glu Leu Ala Phe Thr Arg Pro 1 5 10 15	
55	Tyr Gly Glu Gln Glu Lys Gln Ile Leu Thr Ala Glu Ala Val Glu Phe 20 25 30	
60	Leu Thr Glu Leu Val Thr His Phe Thr Pro Gln Arg Asn Lys Leu Leu 35 40 45	
65	Ala Ala Arg Ile Gln Gln Gln Gln Asp Ile Asp Asn Gly Thr Leu Pro 50 55 60	
70	Asp Phe Ile Ser Glu Thr Ala Ser Ile Arg Asp Ala Asp Trp Lys Ile 65 70 75 80	
75	Arg Gly Ile Pro Ala Asp Leu Glu Asp Arg Arg Val Glu Ile Thr Gly 85 90 95	
80	Pro Val Glu Arg Lys Met Val Ile Asn Ala Leu Asn Ala Asn Val Lys 100 105 110	
85	Val Phe Met Ala Asp Phe Glu Asp Ser Leu Ala Pro Asp Trp Asn Lys 115 120 125	

Val Ile Asp Gly Gln Ile Asn Leu Arg Asp Ala Val Asn Gly Thr Ile
 130 135 140
 5 Ser Tyr Thr Asn Glu Ala Gly Lys Ile Tyr Gln Leu Lys Pro Asn Pro
 145 150 155 160
 Ala Val Leu Ile Cys Arg Val Arg Gly Leu His Leu Pro Glu Lys His
 165 170 175
 10 Val Thr Trp Arg Gly Glu Ala Ile Pro Gly Ser Leu Phe Asp Phe Ala
 180 185 190
 Leu Tyr Phe Phe His Asn Tyr Gln Ala Leu Leu Ala Lys Gly Ser Gly
 195 200 205
 15 Pro Tyr Phe Tyr Leu Pro Lys Thr Gln Ser Trp Gln Glu Ala Ala Trp
 210 215 220
 20 Trp Ser Glu Val Phe Ser Tyr Ala Glu Asp Arg Phe Asn Leu Pro Arg
 225 230 235 240
 Gly Thr Ile Lys Ala Thr Leu Leu Ile Glu Thr Leu Pro Ala Val Phe
 245 250 255
 25 Gln Met Asp Glu Ile Leu His Ala Leu Arg Asp His Ile Val Gly Leu
 260 265 270
 Asn Cys Gly Arg Trp Asp Tyr Ile Phe Ser Tyr Ile Lys Thr Leu Lys
 275 280 285
 30 Asn Tyr Pro Asp Arg Val Leu Pro Asp Arg Gln Ala Val Thr Met Asp
 290 295 300
 35 Lys Pro Phe Leu Asn Ala Tyr Ser Arg Leu Leu Ile Lys Thr Cys His
 305 310 315 320
 Lys Arg Gly Ala Phe Ala Met Gly Gly Met Ala Ala Phe Ile Pro Ser
 325 330 335
 40 Lys Asp Glu Glu His Asn Asn Gln Val Leu Asn Lys Val Lys Ala Asp
 340 345 350
 Lys Ser Leu Glu Ala Asn Asn Gly His Asp Gly Thr Trp Ile Ala His
 355 360 365
 45 Pro Gly Leu Ala Asp Thr Ala Met Ala Val Phe Asn Asp Ile Leu Gly
 370 375 380
 50 Ser Arg Lys Asn Gln Leu Glu Val Met Arg Glu Gln Asp Ala Pro Ile
 385 390 395 400
 Thr Ala Asp Gln Leu Leu Ala Pro Cys Asp Gly Glu Arg Thr Glu Glu
 405 410 415
 55 Gly Met Arg Ala Asn Ile Arg Val Ala Val Gln Tyr Ile Glu Ala Trp
 420 425 430
 Ile Ser Gly Asn Gly Cys Val Pro Ile Tyr Gly Leu Met Glu Asp Ala
 435 440 445
 60 Ala Thr Ala Glu Ile Ser Arg Thr Ser Ile Trp Gln Trp Ile His His
 450 455 460
 65 Gln Lys Thr Leu Ser Asn Gly Lys Pro Val Thr Lys Ala Leu Phe Arg
 465 470 475 480

Gln Met Leu Gly Glu Glu Met Lys Val Ile Ala Ser Glu Leu Gly Glu
 485 490 495
 5 Glu Arg Phe Ser Gln Gly Arg Phe Asp Asp Ala Ala Arg Leu Met Glu
 500 505 510
 Gln Ile Thr Thr Ser Asp Glu Leu Ile Asp Phe Leu Thr Leu Pro Gly
 515 520 525
 10 Tyr Arg Leu Leu Ala
 530
 15 <210> 3
 <211> 20
 <212> DNA
 <213> artificial sequence
 20 <220>
 <221> Primer
 <222> (1)..(20)
 <223> aceB5'-1
 25 <400> 3
 ttcggatcca tgacgaggag 20
 <210> 4
 <211> 20
 <212> DNA
 30 <213> artificial sequence
 <220>
 <221> Primer
 <222> (1)..(20)
 35 <223> aceB5'-2
 <400> 4
 ttgccaacag tgcctgatag 20
 40 <210> 5
 <211> 20
 <212> DNA
 <213> artificial sequence
 45 <220>
 <221> Primer
 <222> (1)..(20)
 <223> aceB3'-1
 50 <400> 5
 atgcttactc acgcctgttg 20
 <210> 6
 <211> 20
 55 <212> DNA
 <213> artificial sequence
 <220>
 <221> Primer
 60 <222> (1)..(20)
 <223> aceB3'-2
 <400> 6
 65 catgtgcaga tgctccatag 20

<210> 7
 <211> 1520
 <212> DNA
 <213> Escherichia coli
 5
 <220>
 <221> misc_feature
 <222> (1)..(1520)
 <223> mutagenic DNA
 10
 <220>
 <221> misc_feature
 <222> (1)..(55)
 <223> technical DNA / residues of polylinker sequence
 15
 <220>
 <221> misc_feature
 <222> (56)..(695)
 <223> parts of the 5' region of the aceB gene and regions lying upstream
 20
 <220>
 <221> misc_feature
 <222> (696)..(758)
 <223> technical DNA / residues of polylinker sequence
 25
 <220>
 <221> misc_feature
 <222> (759)..(1467)
 <223> parts of the 3' region of the aceB gene and regions lying downstream
 30
 <220>
 <221> misc_feature
 <222> (1468)..(1520)
 <223> technical DNA / residues of polylinker sequence
 35
 <400> 7
 ctagatgcat gctcgagcgg cgcgcagtgt gatggatatt tgcagaattc ggcttttcgg 60
 atccatgacg aggagctgca cgatgactga acaggcaaca acaaccgatg aactggcttt 120
 40 cacaaggccg tatggcgagc aggagaagca aattcttact gccgaagcgg tagaatttct 180
 gactgagctg gtgacgcatt ttacgccaca acgcaataaa cttctggcag cgcgcattca 240
 45 gcagcagcaa gatattgata acggaacgtt gcctgatttt atttcggaaa cagcttccat 300
 tcgcatgct gattggaaaa ttcgcgggat tcctgcggac ttagaagacc gccgcgtaga 360
 gataactggc ccggtagagc gcaagatggt gatcaacgcg ctcaacgcca atgtgaaagt 420
 50 ctttatggcc gatttcgaag attcactggc accagactgg aacaaagtga tcgacgggca 480
 aattaacctg cgtgatgcgg ttaacggcac catcagttac accaatgaag caggcaaaat 540
 55 ttaccagctc aagcccaatc cagcggtttt gatttgctcg gtacgcggtc tgcacttgcc 600
 ggaaaaacat gtcacctggc gtggtgaggg aatccccggc agcctgtttg attttgcgct 660
 ctattttctc cacaactatc aggcactggt ggcaaaagcc gaattccagc aactggcg 720
 60 ccgttactag tggatccgag atctgcagaa ttccgcttat gcttactcac gcctgttgat 780
 taaaacctgc cataaacgcg gtgcttttgc gatggcgggc atggcggcgt ttattccgag 840
 65 caaagatgaa gagcacaata accaggtgct caacaaagta aaagcggata aatcgctgga 900

```

    agccaataac ggtcacgatg gcacatggat cgctcaccca ggccttgcgg acacggcaat    960
5   ggcgggtattc aacgacattc tcggctcccg taaaaatcag cttgaagtga tgcgcgaaca    1020
    agacgcgcgcg attactgccg atcagctgct ggcaccttgt gatggtgaac gcaccgaaga    1080
    aggtatgcmc gccaacattc gcgtggctgt gcagtacatc gaagcgtgga tctctggcaa    1140
10  cggctgtgtg ccgatttatg gcctgatgga agatgcggcg acggctgaaa tttcccgtac    1200
    ctcgatctgg cagtggatcc atcatcaaaa aacgttgagc aatggcaaac cggtgaccaa    1260
15  agccttgctc cgccagatgc tgggcgaaga gatgaaagtc attgccagcg aactgggcga    1320
    agaacgttcc tcccaggggc gttttgacga tgccgcacgc ttgatggaac agatcaccac    1380
    ttccgatgag ttaattgatt tcctgaccct gccaggctac cgctgttag cgtaaaccac    1440
20  cacataacta tggagcatct gcacatgaag ccgaattcca gcacactggc ggccgttact    1500
    agtggatccg agctcggtag                                     1520

    <210> 8
25  <211> 1353
    <212> DNA
    <213> Escherichia coli

    <220>
30  <221> misc_feature
    <222> (1)..(1353)
    <223> mutagenic DNA

    <220>
35  <221> misc_feature
    <222> (1)..(3)
    <223> start codon of the deltaaceB allele

    <220>
40  <221> misc_feature
    <222> (1)..(613)
    <223> 5' region of the deltaaceB allele

    <220>
45  <221> misc_feature
    <222> (614)..(676)
    <223> technical DNA /residues of polylinker sequence

    <220>
50  <221> misc_feature
    <222> (677)..(1435)
    <223> 3' region of the deltaaceB allele

    <220>
55  <221> misc_feature
    <222> (1433)..(1435)
    <223> stop codon of the deltaaceB allele

    <400> 8
60  atgactgaac aggcaacaac aaccgatgaa ctggctttca caaggccgta tggcgagcag    60
    gagaagcaaa ttcttactgc cgaagcggta gaatttctga ctgagctggt gacgcatttt    120
    acgccacaac gcaataaact tctggcagcg cgcattcagc agcagcaaga tattgataac    180
65

```

	ggaacggtgc ctgattttat ttcggaaaca gcttccattc gcgatgctga ttggaaaatt	240
	cgcgggattc ctgcggactt agaagaccgc cgcgtagaga taactggccc ggtagagcgc	300
5	aagatggtga tcaacgcgct caacgccaat gtgaaagtct ttatggccga tttcgaagat	360
	tcactggcac cagactggaa caaagtgatc gacgggcaaa ttaacctgcg tgatgcggtt	420
10	aacggcacca tcagttacac caatgaagca ggcaaaattt accagctcaa gcccaatcca	480
	gcggttttga tttgtcgggt acgcggtctg cacttgccgg aaaaacatgt cacctggcgt	540
	ggtgaggcaa tccccggcag cctgtttgat tttgcgtctt atttcttcca caactatcag	600
15	gcactgttgg caaaagccga attccagcac actggcggcc gttactagtg gatccgagat	660
	ctgcagaatt cggtttatgc ttactcacgc ctgttgatta aaacctgcca taaacgcggt	720
20	gcttttgcga tgggcggcat ggcggcgttt attccgagca aagatgaaga gcacaataac	780
	cagggtgctca acaaagtaaa agcggataaa tcgctggaag ccaataacgg tcacgatggc	840
	acatgggatcg ctcacccagg ccttgccggac acggcaatgg cggtattcaa cgacattctc	900
25	ggctcccgtg aaaatcagct tgaagtgatg cgcgacaag acgcgccgat tactgccgat	960
	cagctgctgg caccttgtga tgggtgaacgc accgaagaag gtatgcgcgc caacattcgc	1020
30	gtggctgtgc agtacatcga agcgtggatc tctggcaacg gctgtgtgcc gatttatggc	1080
	ctgatggaag atgccccgac ggctgaaatt tcccgtacct cgatctggca gtggatccat	1140
	catcaaaaaa cgttgagcaa tggcaaaccg gtgaccaaag ccttgttccg ccagatgctg	1200
35	ggcgaagaga tgaaagtcag tgccagcgaa ctgggcgaag aacgtttctc ccaggggcgt	1260
	tttgacgatg ccgcacgctt gatggaacag atcaccactt ccgatgagtt aattgatttc	1320
40	ctgaccctgc caggctaccg cctgttagcg taa	1353

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 January 2003 (30.01.2003)

PCT

(10) International Publication Number
WO 03/008604 A3

(51) International Patent Classification⁷: **C12P 13/06**,
13/08, 13/10, 13/12, 13/14, 13/20, 13/22, 13/24

(21) International Application Number: **PCT/EP02/07352**

(22) International Filing Date: **3 July 2002 (03.07.2002)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
101 35 051.1 18 July 2001 (18.07.2001) DE
60/306,867 23 July 2001 (23.07.2001) US

(71) Applicant (for all designated States except US): **DE-GUSSA AG [DE/DE]**; Bennigsenplatz 1, 40474 Düsseldorf (DE).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **HERMANN, Thomas [DE/DE]**; Zirkonstrasse 8, 33739 Bielefeld (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:


— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report

[Continued on next page]

(54) Title: **PROCESS FOR THE PREPARATION OF L-AMINO ACIDS USING STRAINS OF THE ENTEROBACTERIACEAE FAMILY WHICH CONTAIN AN ATTENUATED ACEB GENE**



(57) Abstract: The invention relates to a process for the preparation of L-amino acids, in particular L-threonine, in which the following steps are carried out: a) fermentation of microorganisms of the Enterobacteriaceae family which produce the desired L-amino acid and in which the aceB gene, or the nucleotide sequence which codes for this, is attenuated, in particular eliminated, b) concentration of the L-amino acid in the medium or in the cells of the bacteria, and c) isolation of the L-amino acid.

WO 03/008604 A3



(88) Date of publication of the international search report:
4 December 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/07352

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12P13/06 C12P13/08 C12P13/10 C12P13/12 C12P13/14
C12P13/20 C12P13/22 C12P13/24

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 378 616 A (TUJIMOTO NOBUHARU ET AL) 3 January 1995 (1995-01-03) page 2, column 2, paragraph 2	1-7
A	WO 94 28154 A (NUTRASWEET CO) 8 December 1994 (1994-12-08) page 1, line 10 - line 14; table 2	1-7
A	CHUNG T ET AL: "Glyoxylate bypass operon of Escherichia coli: cloning and determination of the functional map." JOURNAL OF BACTERIOLOGY. US, vol. 170, no. 1, January 1988 (1988-01), pages 386-392, XP008015355 ISSN: 0021-9193 the whole document	1-7



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z document member of the same patent family

Date of the actual completion of the international search

3 April 2003

Date of mailing of the international search report

17/04/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kools, P

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/07352

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 347 318 A (MIWA KIYOSHI ET AL) 31 August 1982 (1982-08-31) the whole document	1-7
A	REINSCHIED D J ET AL: "Malate synthase from <i>Corynebacterium glutamicum</i> : sequence analysis of the gene and biochemical characterization of the enzyme" MICROBIOLOGY, vol. 140, no. 11, 1994, pages 3099-3108, XP008012735 Reading, GB ISSN: 1350-0872 the whole document	1-7
A	LANDGRAF J R ET AL: "THE ROLE OF H-NS IN ONE CARBON METABOLISM" BIOCHIMIE, MASSON, PARIS, FR, vol. 76, no. 10/11, 1994, pages 1063-1070, XP008014239 ISSN: 0300-9084 the whole document	6
A	RAE J L ET AL: "Sequences and expression of pyruvate dehydrogenase genes from <i>Pseudomonas aeruginosa</i> ." JOURNAL OF BACTERIOLOGY, vol. 179, no. 11, June 1997 (1997-06), pages 3561-3571, XP002237063 US ISSN: 0021-9193 abstract	1-7
E	WO 03 008600 A (DEGUSSA ;HERMANN THOMAS (DE)) 30 January 2003 (2003-01-30) claim 7	7
E	WO 03 008602 A (DEGUSSA ;HERMANN THOMAS (DE)) 30 January 2003 (2003-01-30) claim 7	7
E	WO 03 008603 A (DEGUSSA ;HERMANN THOMAS (DE)) 30 January 2003 (2003-01-30) claim 7	7
E	WO 03 008616 A (DEGUSSA ;HERMANN THOMAS (DE)) 30 January 2003 (2003-01-30) claim 7	7

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/07352

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5378616	A	03-01-1995	BR 9203053 A FR 2680178 A1 JP 3106714 B2 JP 5244970 A PH 30131 A US 5393671 A	30-03-1993 12-02-1993 06-11-2000 24-09-1993 21-01-1997 28-02-1995
WO 9428154	A	08-12-1994	AU 6960894 A CA 2140527 A1 EP 0660875 A1 JP 7509375 T WO 9428154 A1	20-12-1994 08-12-1994 05-07-1995 19-10-1995 08-12-1994
US 4347318	A	31-08-1982	JP 1029559 B JP 1552063 C JP 55131397 A DE 3012921 A1 FR 2453216 A1 GB 2049670 A , B	12-06-1989 23-03-1990 13-10-1980 23-10-1980 31-10-1980 31-12-1980
WO 03008600	A	30-01-2003	DE 10135051 A1 WO 03008600 A2 WO 03008602 A2 WO 03008603 A2 WO 03008604 A2 WO 03008616 A2	06-02-2003 30-01-2003 30-01-2003 30-01-2003 30-01-2003 30-01-2003
WO 03008602	A	30-01-2003	DE 10135051 A1 WO 03008600 A2 WO 03008602 A2 WO 03008603 A2 WO 03008604 A2 WO 03008616 A2	06-02-2003 30-01-2003 30-01-2003 30-01-2003 30-01-2003 30-01-2003
WO 03008603	A	30-01-2003	DE 10135051 A1 WO 03008600 A2 WO 03008602 A2 WO 03008603 A2 WO 03008604 A2 WO 03008616 A2	06-02-2003 30-01-2003 30-01-2003 30-01-2003 30-01-2003 30-01-2003
WO 03008616	A	30-01-2003	DE 10135051 A1 WO 03008600 A2 WO 03008602 A2 WO 03008603 A2 WO 03008604 A2 WO 03008616 A2	06-02-2003 30-01-2003 30-01-2003 30-01-2003 30-01-2003 30-01-2003